
Antenatal screening



All infants of HBsAg-positive mothers should have HBIg within 24 hours of birth, as well as the hepatitis vaccine.

* Terrault NA, et al. Hepatology, vol 67, no. 4, 2018 Lampertico P, et al. Journal of Hepatology. 2017, vol 67, 370-398.



The Hepatitis Foundation of New Zealand

PO Box 647, Whakatane 3158

Phone: 0800 33 20 10 Fax: (07) 307 1266

Email: hepteam@hfnz.nz

Web: www.hepatitisfoundation.org.nz

C8.4 Antenatal Screening		
Issue Date: September 2020	Version 2	Next Review Date: September 2022
DOCUMENTATION – The electronic version is the most up to date version. The Hepatitis Foundation accepts no responsibility for		
the consequences that arise from using out of date printed copies of this document		

Document Control Version History

Version	Date Issued	Brief Summary of Change	Owner's Name
Version 1	September 2020	New document	Alex Lampen-Smith
Version 2	September 2021	Review	Alex Lampen-Smith

Review and consultation	Next annual review: September 2022
process:	CEO to oversee process.
Responsibility for	Day to day responsibility for implementation: CEO
Implementation:	

Revisions:		
Date	Author	Description
September 2021	Alex Lampen-Smith	Annual review

Distribution	Communication and distribute the policy by Internal email,
Methods:	electronic copy held on shared drive.

C8.4 Antenatal Screening		
Issue Date: September 2020	Version 2	Next Review Date: September 2022
DOCUMENTATION – The electronic version is the most up to date version. The Hepatitis Foundation accepts no responsibility for		

DOCUMENTATION – The electronic version is the most up to date version. The Hepatitis Foundation accepts no responsibility for the consequences that arise from using out of date printed copies of this document